



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

DESCRIPTION OF A NEW SPECIES OF *PIPILO*
FROM MOUNT ORIZABA, MEXICO.

BY ULYSSES O. COX.

While on Mount Orizaba, Mexico, as a member of Dr. Scovell's party during the summer of 1891 I made a small collection of birds of the region which Mr. Robert Ridgway of the U. S. National Museum has had the kindness to identify for me. Among them Mr. Ridgway finds a *Pipilo* which appears to be a new species. I have given it the specific name *orizabæ* and append here Mr. Ridgway's description and comparison with related species.

Pipilo orizabæ Cox, sp. nov.

"SP. CHAR.—Most like *P. maculatus* Sw., but no black whatever on upper parts, which are plain grayish brown; white markings of scapulars and wing feathers nearly obsolete, sides and flanks much paler in color, and size somewhat greater.

"*Adult male* (No. 132,720, U. S. National Museum, Mount Orizaba, Mexico, alt. 11,000 feet; Ulysses O. Cox): Above uniform dull grayish brown, slightly tinged with olive, scarcely darker on the head; outermost scapulars with marginal elongated spots of white, and lowermost middle and greater wing-coverts with more roundish terminal spots of the same; no white markings on remiges, but primaries edged with light brownish gray; tail (except middle feathers) very dark brown, the three outermost feathers with a large, abruptly defined, white spot terminating inner webs, that on the lateral feather nearly 1 inch long. Chin, throat, and chest blackish brown, gradually lightening on sides of head and grading gradually into color of crown, the throat spotted with white beneath the surface; sides, flanks, and under tail-coverts ochraceous-buff, somewhat deeper anteriorly; median lower parts white, broadest and purest anteriorly, next to blackish brown of the chest. Bill black; legs light brown, toes darker. Total length (skin), 8.30¹; wing, 3.35; tail, 3.85; exposed culmen, 0.60; tarsus, 1.17; middle toe, 0.78.

"This bird is obviously quite distinct from both *P. maculatus* and *P. macronyx*, as well as their hitherto described allies. The pure white and abruptly defined tail-spots, and entire

¹ Before skinning, 21 cm., = about 8.27 inches.

absence of green or yellow from any part of the plumage separate it at once from *P. macronyx*, *P. virescens* and *P. complexus*; the entire absence of black from the upper parts from *P. maculatus* and *P. submaculatus*, while it differs from all in having the entire top and sides of the head grayish brown, like the back, instead of black, like the chest. The general color of the upper parts is very much that of the darker examples of *P. fuscus*, but the color has, in certain lights, a very preceptible olivaceous cast."

BIRDS OF ONEIDA COUNTY, NEW YORK.

BY EGBERT BAGG.

SINCE the publication of the article by Dr. Ralph and myself in 'The Auk' for July, 1890, I have been able to add the following to our local list. These eight new records, together with *Vireo solitarius plumbeus*, recorded from the adjoining county of Madison, by Gerrit S. Miller, Jr., in the last number of this journal, make the total number of species and subspecies recorded from "Oneida County, N. Y., and its immediate vicinity" two hundred and thirty-nine.

***Rissa tridactyla*.**—A young male of this species was killed at Constantia, Oswego Co. (on Oneida Lake), Nov. 9, 1890, by Robert J. Hughes.

***Rynchops nigra*.**—I have lately examined a specimen of this bird in a taxidermist's shop in Utica, which was brought in in the fall of 1893 (October?) and reported to have been killed near Whitesboro. Baird, Brewer, and Ridgway say of this species: "It is never known to be driven astray by any storm, however violent."

***Æstrelata hasitata*.**—A male of this rare straggler was shot at Verona Beach, on Oneida Lake, August 28, 1893, by the Rev. G. A. Biederman, who presented it to Alex. H. Moore, a young ornithologist of Utica, who mounted and preserved it. Mr. B. reports that there were two birds together, but careful search shortly afterward failed to find the other, and it may have been some other species. Through the courtesy of Mr. Moore, who brought the bird to me for identification while in the flesh,